

---

## Short Database Report

# KRITI – the vegetation of Crete database

Erwin Bergmeier

**Abstract:** The TURBOVEG database on the Cretan covers the South Aegean island of Crete including surrounding islands. It contains plot data on all kinds of woody and non-woody vascular plant vegetation. Bryophytes are included in exceptional cases. It includes relevés from all available sources but chiefly own unpublished relevés. The database provides information on about 90% of the regional flora and of more than 100 endemics. KRITI (GIVD ID EU-GR-001) is a database to be continued. Publications dealing with selected vegetation types or areas have been published.

**Keywords:** Aegean; Greece; Mediterranean vegetation.

GIVD Database ID: EU-GR-001	Last update: 2011-07-06
<b>KRITI</b>	
<b>Scope:</b> Plot data (almost exclusively phytosociological relevés) of all kinds of vegetation from the island of Crete and surrounding islands (Greece).	
<b>Status:</b> completed and continuing	<b>Period:</b> 1945-2010
<b>Database manager(s):</b> Erwin Bergmeier (erwin.bergmeier@bio.uni-goettingen.de)	
<b>Owner:</b> Erwin Bergmeier (private)	
<b>Web address:</b> [NA]	
<b>Availability:</b> according to a specific agreement	<b>Online upload:</b> no
<b>Database format(s):</b> TURBOVEG	<b>Export format(s):</b> [NA]
<b>Publication:</b> Bergmeier, E. 2001: Von der pflanzensoziologischen Aufnahme zur Synopsis: eine Vegetationsdatenbank von Kreta. - Berichte der Reinhold-Tüxen-Gesellschaft 13: 71-82.	
<b>Plot type(s):</b> normal plots	<b>Plot-size range:</b> 1-400 m <sup>2</sup>
<b>Non-overlapping plots:</b> 6,500	<b>Estimate of existing plots:</b> 8,000
<b>Total plot observations:</b> 6,500	<b>Number of sources:</b> 0
<b>Countries:</b> GR: 100.0%	<b>Completeness:</b> 81%
<b>Forest:</b> [NA] — <b>Non-forest:</b> [NA]	<b>Valid taxa:</b> 1,600
<b>Guilds:</b> all vascular plants: 100%; bryophytes (terricolous or aquatic): 10%; lichens (terricolous or aquatic): 1%; algae (terricolous or aquatic): 0%	
<b>Environmental data:</b> altitude: 100%; slope aspect: 100%; slope inclination: 100%; microrelief: 35%; soil depth: 100%; surface cover other than plants (open soil, litter, bare rock etc.): 25%; other soil attributes: 75%	
<b>Performance measure(s):</b> cover: 100%	
<b>Geographic localisation:</b> GPS coordinates (precision 25 m or less): 75%; point coordinates less precise than GPS, up to 1 km: 5%; small grid (not coarser than 10 km): 20%	
<b>Sampling periods:</b> 1940-1949: 1.0%; 1960-1969: 1.0%; 1970-1979: 3.0%; 1980-1989: 6.0%; 1990-1999: 30.0%; 2000-2009: 55.0%; 2010-2019: 4.0%	
<b><i>Information as of 2012-07-12; further details and future updates available from <a href="http://www.givd.info/ID/EU-GR-001">http://www.givd.info/ID/EU-GR-001</a></i></b>	

Erwin Bergmeier (erwin.bergmeier@bio.uni-goettingen.de)  
Vegetation Analysis & Phytodiversity, University of Göttingen, Untere Karspüle 2, 37073 Göttingen, GERMANY