

Fairy Circles of the Namib Desert - Ecosystem engineering by subterranean social insects

Author: Jürgens, N.

Published by: Biodiversity & Ecology, 7, 376 pp

Klaus Hess Publishers, Göttingen & Windhoek

DOI: [10.7809/b-e.vol_07](https://doi.org/10.7809/b-e.vol_07)

Fairy Circles of the Namib Desert

Ecosystem engineering by subterranean social insects

by

Norbert Jürgens

with contributions from

**Alicia Geppert
Alexander Gröngröft
Felicitas Gunter
Joh R. Henschel
Katharina J. Huber
Joe McAuliffe
Jens Oldeland
Jörg Overmann
Javier Pascual
Mike Picker
Rasmus Revermann
Priscilla Sichone
Johannes Sikorski
Andrey Yurkov**

Klaus Hess Publishers
Göttingen & Windhoek

2022

Abstract

On the eastern edge of the Namib Desert—probably the oldest continuous desert in the world—millions of ochre-coloured bare patches in the grass have created landscapes of unique beauty that are often regarded as one of the greatest mysteries in ecology: the fairy circles of the Namib Desert.

Building on more than four decades of ecological research in all parts of the Namib in Angola, Namibia, and South Africa, the authors here present the natural history of the phenomenon, with all the facts understood so far, and also describe the remaining open questions.

In the Namib Desert, sand termites from soil-dwelling termite families have been found to cause the fairy circles. These termites must be understood as ecosystem engineers that—comparable to the beavers in rivers of the northern hemisphere—profoundly alter the landscape and make permanent life in the desert possible.

The termites create the bare spots in the vegetation through localized herbivory, with the ecological effect that subsequent rainfall is not immediately absorbed and transpired by plants. Instead, the sandy soils in the open spots allow rapid percolation of the rainwater to soil depths below 50 cm, where water remains stored even during prolonged periods of drought. This water storage enables the termites to live permanently even in dry climatic zones with less than 100 mm of annual precipitation. At the same time, numerous other animals and plants benefit from these underground water resources.

Although water and fresh plant biomass are the main attraction, several animals also use the termites as a source of food. The most notable predator is the aardvark, but the list also includes several carnivorous and omnivorous ants, which are eaten in turn by spiders, ant lions, reptiles, birds, and many other members of a diverse food web.

The termite nest within the fairy circles contains a special microbiome on its own, which involves various bacteria and fungi some of which produce organic and inorganic gases such as methane.

A comparative look beyond the Namib Fairy Circles shows that also other social insects are important and often underrated ecosystem engineers and produce similar phenomena within the Namib Desert and in other climates.

However, the fairy circles of the Namib Desert are probably the most astounding and efficient product of adaptation to extreme aridity. The sand termites manipulate the local topsoil hydrology and thereby turn an ephemeral desert into a permanently habitable grassland.

Bibliography

- Abdullah, W., Elsayed, W.M., Abdelshafeek, K.A., Nazif, N.M. & Singab, A.N.B. (2016) Chemical constituents and biological activities of *Cleome* genus: a brief review. *International Journal of Pharmacognosy and phytochemical research*, **8**, 777–787.
- Aho, K. A. (2013) *Foundational and applied statistics for biologists using R*. CRC Press.
- Aho, K.A. (2018) Asbio: A Collection of Statistical Tools for Biologists. R package version 1.5-3. [cited 08 November 2018].
- Aho, K.A. (2022) Asbio: A Collection of Statistical Tools for Biologists. R package version 1.8-2.
- Albrecht, C.F., Joubert, J.J. & De Rycke, P.H. (2001) Origin of the enigmatic, circular, barren patches ('Fairy Rings') of the pro-Namib. *South African Journal of Science*, **97**, 23–27.
- António, T.E.M. (2022) Plant diversity and vegetation patterns in the north of the Kaokoland endemic centre, Iona National Park. Doctoral dissertation, Namibia University of Science and Technology, Namibia.
- Apel, C. (2018) Spatial and activity patterns of *Ocymyrmex velox* in the Marienfluss Valley, Namibia. Bachelor thesis, Institute of Plant Science and Microbiology, University of Hamburg.
- Arnqvist, G. & Mårtensson, T. (1998) Measurement error in geometric morphometrics: Empirical strategies to assess and reduce its impact on measures of shape. *Acta Zoologica Academiae Scientiarum Hungaricae*, **44**, 73–96.
- Arshad, M.A. (1982) Influence of the termite *Macrotermes michaelensi* (Sjost) on soil fertility and vegetation in a semi-arid savannah ecosystem. *Agro-ecosystems*, **8**, 47–58.
- Austin, J.W., Bagnères, A.G., Szalanski, A.L., Scheffrahn, R.H., Heintschel, B.P., Messenger, M.T., Clément, J.-L. & Gold, R. E. (2007) *Reticulitermes mallei* (Isoptera: Rhinotermitidae): A valid Nearctic subterranean termite from Eastern North America. *Zootaxa*, **26**, 1–26.
- Barbosa, L.A.G. (1970) *Carta Fitogeográfica da Angola*. Instituto de Investigação Científica de Angola, Luanda.
- Bardunias, P.M., Calovi, D.S., Carey, N., Soar, R., Turner, J.S., Nagpal, R. & Werfel, J. (2020) The extension of internal humidity levels beyond the soil surface facilitates mound expansion in *Macrotermes*. *Proceedings of the Royal Society B: Biological Sciences*, **287**, 20200894.
- Becker, T. (2007) The phenomenon of fairy circles in Kaokoland (NW Namibia). *Basic and Applied Dryland Research*, **1**, 121–137.
- Becker, T. & Getzin, S. (2000) The fairy circles of Kaokoland (North-West Namibia) – origin, distribution, and characteristics. *Basic and Applied Ecology*, **1**, 149–159.
- Bischoff, G. (2013) Activity Patterns and Behaviour of *Tetramorium* Ants in Fairy Circle Ecosystems. Bachelor thesis, Institute of Plant Science and Microbiology, University of Hamburg.
- Bohn, C. (2014) Population genetics on the sand termite *Psammotermes allocerus* in Namib Desert fairy circles. Master thesis, Institute of Plant Science and Microbiology, University of Hamburg.
- Bollig, M. (1997) Risk and risk minimisation among Himba pastoralists in northwestern Namibia. *Nomadic Peoples*, **1**, 66–89.
- Bollig, M., & Gewald, J.B. (2000) People, Cattle and Land. *Transformations of a Pastoral Society in Southwestern Africa*. Köppé, Köln.
- Bonanomi, G., Incerti, G., Stinca, A., Carteni, F., Giannino, F. & Mazzoleni, S. (2014) Ring formation in clonal plants. *Community Ecology*, **15**, 77–86.
- Bonanomi, G., Mingo, A., Incerti, G., Mazzoleni, S. & Allegrezza, M. (2012) Fairy rings caused by a killer fungus foster plant diversity in species-rich grassland. *Journal of Vegetation Science*, **23**, 236–248.
- Bothma, J.d.P. & Nel, J.A.J. (1980) Winter food and foraging behaviour of the aardwolf *Proteles cristatus* in the Namib-Naukluft Park. *Madoqua*, **12**, 141–149.
- Bourguignon, T., & Roisin, Y. (2011). Revision of the termite family Rhinotermitidae (Isoptera) in New Guinea. *ZooKeys*, **148**, 55–103.
- Bourguignon, T., Šobotník, J., Sillam-Dussès, D., Jiroš, P., Hanus, R., Roisin, Y. & Miura, T. (2012) Developmental Pathways of *Psammotermes hybostoma* (Isoptera: Rhinotermitidae): Old Pseudergates Make up a New Sterile Caste. *PLoS One*, **7**, e44527.
- Brandl, R., Hacker, M., Epplen, J.T. & Kaib, M. (2005) High gene flow between populations of *Macrotermes michaelensi* (Isoptera, Termitidae). *Insectes Sociaux*, **52**, 344–349.
- Breiman, L., Friedman, J., Stone, C.J. & Olshen, R.A. (1984) *Classification and regression trees* - crc press. Boca Raton, Florida.
- Brown, G., Scherber, C., Ramos Jr, P. & Ebrahim, E.K. (2012) The effects of harvester ant (*Messor ebeninus* Forel) nests on vegetation and soil properties in a desert dwarf shrub community in north-eastern Arabia. *Flora-Morphology, Distribution, Functional Ecology of Plants*, **207**, 503–511.
- Brune, A. (2014) Symbiotic digestion of lignocellulose in termite guts. *Nature Reviews Microbiology*, **12**, 168–180.
- Burke, A. (2006) *The Sperrgebiet: Managing its biodiversity*. EnviroScience and Namibia Nature Foundation, Oranjemund and Windhoek, 100 pages.
- Burke, A. & Loots, S. (2020) Plant endemics of the TsauKhaeb (Sperrgebiet) National Park. *Namibian Journal of Environment*, **4**, 62–70.
- Caldwell, M.M., Dawson, T.E. & Richards, J.H. (1998) Hydraulic lift: consequences of water efflux from the roots of plants. *Oecologia*, **113**, 151–161.
- Campbell, N.A. & Atchley, W.R. (1981) The geometry of canonical variate analysis. *Systematic Biology*, **30**, 268–280.
- Carrington, D. (2014) Namibia's 'fairy circles': Nature's greatest mystery? CNN, <https://edition.cnn.com/2014/06/13/world/africa/namibia-fairy-circles-nature-greatest-mystery/index.html>.
- Carteni, F., Marasco, A., Bonanomi, G., Mazzoleni, S., Rietkerk, M. & Giannino, F. (2012) Negative plant soil feedback explaining ring formation in clonal plants. *Journal of Theoretical Biology*, **313**, 153–161.
- Chouvenç, T., Šobotník, J., Engel, M. S., & Bourguignon, T. (2021). Termite evolution: mutualistic associations, key innovations, and the rise of Termitidae. *Cellular and Molecular Life Sciences*, **78**(6), 2749–2769.
- Clark, P.J. & Evans, F.C. (1954) Distance to nearest neighbor as a measure of spatial relationships in populations. *Ecology*, **35**, 445–453.
- Coaton, W.G.H. (1958) The hodotermitid harvester termites of South Africa. Department of Agriculture, Union of South Africa.
- Coaton, W.G.H. (1962) Control of the hodotermitid harvester termites in the Karoo. *Journal of the Entomological Society of Southern Africa*, **25**, 318–327.
- Coaton, W.G.H. (1981) Fossilised nests of Hodotermitidae (Isoptera) from the Clanwilliam District, Cape Province. *Journal of the Entomological Society of Southern Africa*, **44**, 79–81.
- Coaton W.G.H. & Sheasby J.L. (1972) Preliminary report on a survey of the termites (Isoptera) of South West Africa. *Cimbebasia Memoirs* **2**, 1–129.
- Coaton W.G.H. & Sheasby J.L. (1973) National Survey of the Isoptera of Southern Africa. 3. The Genus *Psammotermes* Desneux (Rhinotermitidae). *Cimbebasia*, **3**, 19–28.
- Coaton, W.G.H. & Sheasby, J.L. (1973) National Survey of the Isoptera of southern Africa. 1. The genus *Baucaliotermes* Sands (Termitidae: Nasutitermitinae). *Cimbebasia*, **3**, 2–7.
- Coaton, W.G.H. & Sheasby, J.L. (1974) National survey of the Isoptera of southern Africa. 6. The Genus *Microhodotermes* Sjöstedt (Hodotermitidae). *Cimbebasia*, **3**, 47–59.
- Coaton, W.G.H. & Sheasby, J.L. (1974) National survey of the Isoptera of southern Africa. 6. *Hodotermes mossambicus* (Hagen) *Cimbebasia*, **3**, 106–137.

- Coaton, W.G.H. & Sheasby, J.L. (1975) National Survey of the Isoptera of Southern Africa. 10. The Genus *Hodotermes* Hagen (Hodotermitidae). *Cimbebasia*, **3**, 105–138.
- Coaton, W.G.H. & Sheasby, J.L. (1976) National survey of the Isoptera of southern Africa. 11. The genus *Coptotermes* Wasmann (Rhinotermitidae: Coptotermitinae). *Cimbebasia*, **3**, 140–172.
- Coetzee, M.E. (2021) Soils of the Sceleton Coast National Park and SCIONA project area in Namibia. Windhoek, Namibia.
- Cohen, J. (1960) A Coefficient of Agreement for Nominal Scales. *Educational and Psychological Measurement*, **20**, 37–46.
- Cox, G.W. (1987) The origin of vegetation circles on stony soils of the Namib Desert near Gobabeb, South West Africa/Namibia. *Journal of Arid Environments*, **13**, 237–244.
- Cox, G.W., Lovegrove, B.G. & Siegfried, W.R. (1987) The small stone content of mima-like mounds in the South African Cape region: implications for mound origin. *Catena*, **14**, 165–176.
- Cramer, M.D. & Barger, N.N. (2013) Are Namibian “fairy circles” the consequence of self-organizing spatial vegetation patterning? *PloS one*, **8**, e70876.
- Cramer, M.D., Barger, N.N. & Tschinkel, W.R. (2017) Edaphic properties enable facilitative and competitive interactions resulting in fairy circle formation. *Ecography*, **40**, 1210–1220.
- Cramer, M.D. & Midgley, J.J. (2015) The distribution and spatial patterning of mima-like mounds in South Africa suggests genesis through vegetation induced aeolian sediment deposition. *Journal of Arid Environments*, **119**, 16–26.
- Curtis, B.A. (1985) The dietary spectrum of the Namib Desert dune ant *Camponotus detritus*. *Insectes Sociaux*, **32**, 78–85.
- d'Herbès, J.M., Valentin, C., Tongway, D.J. & Leprun, J.C. (2001) Banded vegetation patterns and related structures. *Banded vegetation patterning in arid and semiarid environments* (ed. by P.J. Edwards, R.M. May and N.R. Webb), pp. 1–19. Springer, New York.
- Dangerfield, J.M., McCarthy, T.S. & Ellerly, W.N. (1998) The mound-building termite *Macrotermes michaelseni* as an ecosystem engineer. *Journal of Tropical Ecology*, **14**, 507–520.
- Danin, A. & Orshan, G. (1995) Circular arrangement of *Stipagrostis ciliata* clumps in the Negev, Israel and near Gokae, Namibia. *Journal of Arid Environments*, **30**, 307–313.
- Darimont, C.T., Reimchen, T.E. & Paquet, P.C. (2003) Foraging behaviour by gray wolves on salmon streams in coastal British Columbia. *Canadian Journal of Zoology*, **81**, 349–353.
- Dayrat, B. (2005) Towards integrative taxonomy. *Biological Journal of the Linnean Society*, **85**, 407–415.
- Deblauwe, V., Barbier, N., Couteron, P., Lejeune, O. & Bogaert, J. (2008) The global biogeography of semi-arid periodic vegetation patterns. *Global Ecology and Biogeography*, **17**, 715–723.
- Deblauwe, V., Couteron, P., Bogaert, J. & Barbier, N. (2012) Determinants and dynamics of banded vegetation pattern migration in arid climates. *Ecological Monographs*, **82**, 3–21.
- Dinter, K., Paarmann, W., Peschke, K. & Arndt, E. (2002) Ecological, behavioural and chemical adaptations to ant predation in species of *Thermophilum* and *Graphipterus* (Coleoptera: Carabidae) in the Sahara Desert. *Journal of Arid Environments*, **50**, 267–286.
- Duncan, F.D. & Hewitt, P.H. (2009) Observations on the foraging behaviour of the harvester termite, *Hodotermes mossambicus* (Hagen) (Isoptera: Hodotermitidae). *Bulletin of Entomological Research*, **79**, 631–642.
- Dunkerley, D. (2018) Banded vegetation in some Australian semi-arid landscapes: 20 years of field observations to support the development and evaluation of numerical models of vegetation pattern evolution. *Desert*, **23**, 165–187.
- Eggleton, P. (2010) An introduction to termites: biology, taxonomy and functional morphology. *Biology of Termites: A Modern Synthesis* (ed. by D.E. Bignell, Y. Roisin, and N. Lo), pp. 1–26. Dordrecht, Springer.
- Eicker, A., Theron, G.K. & Grobbelaar, N. (1982) 'n Mikrobiologiese studie van 'kaal kolle' in die Giribesvlakte van Kaokoland, SWA-Namibië. *South African Journal of Botany*, **1**, 69–74.
- Eldridge, D.J. (1994) Nests of ants and termites influence infiltration in a semi-arid woodland. *Pedobiologia*, **38**, 481–492.
- ESRI (2011) ArcGIS Desktop: Release 10. *Environmental Systems Research Institute*, CA.
- Feddes, R.A., Hoff, H., Bruen, M., Dawson, T., De Rosnay, P., Dirmyer, P., Jackson, R.B., Kabat, P., Kleidon, A. & Lilly, A. (2001) Modeling root water uptake in hydrological and climate models. *Bulletin of the American Meteorological Society*, **82**, 2797–2810.
- Fick, S.E. & Hijmans, R.J. (2017) WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology*, **37**, 4302–4315.
- Fox-Dobbs, K., Doak, D.F., Brody, A.K. & Palmer, T.M. (2010) Termites create spatial structure and govern ecosystem function by affecting N₂ fixation in an East African savanna. *Ecology*, **91**, 1296–1307.
- Galt, N.S. & Meyer, J.J.M. (2018) The southern Namibian (Garub) fairy circles: The role of the phytotoxic properties of the toxic succulent, *Euphorbia gummifera*. *South African Journal of Botany*, **115**, 286.
- Geerts, S., Van der Linden, J. & Van der Linden, E. (2016) The ecology and foraging behaviour of the harvester termite, *Baicaliotermes hainesi* in semi-arid grasslands in the northwestern interior of South Africa. *Insects Sociaux*, **63**, 457–465.
- Getzin, S., Nambwandja, A., Holch, S. & Wiegand, K. (2021) Revisiting Theron's hypothesis on the origin of fairy circles after four decades: *Euphorbias* are not the cause. *BMC ecology and evolution*, **21**, 1–23.
- Getzin, S., Wiegand, K., Wiegand, T., Yizhaq, H., Von Hardenberg, J. & Meron, E. (2015a) Adopting a spatially explicit perspective to study the mysterious fairy circles of Namibia. *Ecography*, **38**, 1–11.
- Getzin, S., Wiegand, K., Wiegand, T., Yizhaq, H., Von Hardenberg, J. & Meron, E. (2015b) Clarifying misunderstandings regarding vegetation self-organization and spatial patterns of fairy circles in Namibia: a response to recent termite hypotheses. *Ecological Entomology*, **40**, 669–675.
- Getzin, S. & Yizhaq, H. (2019) Unusual Namibian fairy circle patterns in heterogeneous and atypical environments. *Journal of Arid Environments*, **164**, 85–89.
- Getzin, S., Yizhaq, H., Bell, B., Erickson, T.E., Postle, A.C., Katra, I., Tzuk, O., Zelnik, Y.R., Wiegand, K., Wiegand, T. & Meron, E. (2016) Discovery of fairy circles in Australia supports self-organization theory. *Proceedings of the National Academy of Sciences*, **113**, 3551–3556.
- Getzin, S., Yizhaq, H., Cramer, M.D. & Tschinkel, W.R. (2019) Contrasting global patterns of spatially periodic fairy circles and regular insect nests in drylands. *Journal of Geophysical Research-Biogeosciences*, **124**, 3327–3342.
- Getzin, S., Yizhaq, H. & Tschinkel, W.R. (2021) Definition of “fairy circles” and how they differ from other common vegetation gaps and plant rings. *Journal of Vegetation Science*, **32**, e13092.
- Giess, W. (1971) A preliminary vegetation map of South West Africa. *Dinteria*, **4**, 5–114.
- Gilad, E., von Hardenberg, J., Provenzale, A., Shachak, M., & Meron, E. (2004) Ecosystem engineers: from pattern formation to habitat creation. *Physical Review Letters*, **93**, 098105.
- Gilad, E., von Hardenberg, J., Provenzale, A., Shachak, M., & Meron, E. (2007) A mathematical model of plants as ecosystem engineers. *Journal of Theoretical Biology*, **244**, 680–691.
- Gossweiler J. & Mendonça, F.A. (1939) Carta Fitogeográfica de Angola. Ministério das Colônias, Lisboa.
- Goudie, A.S. (1972) Climate, weathering, crust formation, dunes, and fluvial features of the Central Namib Desert, near Gobabeb, South West Africa. *Madoqua*, **2** (1), 54–62.
- Goudie, A.S. (2022) Nebkhas: An essay in aeolian biogeomorphology. *Aeolian Research*, **54**, 100772.
- Goudie, A.S. & Viles, H. (2014) *Landscape and Landforms of Namibia*. Springer Dordrecht.

- Govorushko, S. (2019) Economic and ecological importance of termites: A global review. *Entomological Science*, **22**, 21–35. <https://doi.org/10.1111/ens.12328>
- Grabovsky, V.I. (2018) A model of vegetation cover in conditions of resource scarcity: Fairy rings in Namibia. *Biology Bulletin Reviews*, **8**, 169–180.
- Gregory, P.H. (1982) Fairy rings; free and tethered. *Bulletin of the British Mycological Society*, **16**, 161–163.
- Grohmann, C. (2010) Termite mediated heterogeneity of soil and vegetation patterns in a semi-arid savanna ecosystem in Namibia. Doctoral dissertation, Universität Würzburg.
- Grohmann, C., Oldeland, J., Stoyan, D. & Linsenmair, K.E. (2010) Multi-scale pattern analysis of a mound-building termite species. *Insectes Sociaux*, **57**, 477–486.
- Grube, S. (1993) Untersuchungen zum Wasser-gleichgewicht bei Termiten in ariden Lebensräumen am Beispiel von *Psammotermes allocerus* Silvestri. Doctoral dissertation, Freie Universität Berlin.
- Grube, S. (2000) Soil dumps: indicators of foraging activity by *Hodotermes mossambicus* (Hagen) (Isoptera: hodotermitidae) in northern Namibia? *Cimbebasia*, **16**, 269–270.
- Grube, S. (2002) The fairy circles of Kaokoland (Northwest Namibia) – is the harvester termite *Hodotermes mossambicus* the prime causal factor in circle formation? *Basic and Applied Ecology*, **3**, 367–370.
- Grube, S. & Rudolph, D. (1995) Termites in arid environments: the water balance of *Psammotermes allocerus* Silvestri. *Mitteilungen der Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie*, **10**, 665–668.
- Gunter, F., Oldeland, J., Picker, M.D., Henschel, J.R. & Jürgens, N. (2022) Cryptic subterranean diversity: Regional phylogeography of the sand termite *Psammotermes allocerus* Silvestri, 1908 in the wider Namib region. *Organisms Diversity & Evolution*, **1**–12. <https://doi.org/10.1007/s13127-022-00580-w>
- Gutteridge, G. (2017) Invertebrates of Southern Africa and their Tracks & Signs. Jacana Media, Auckland Park. 409 pp.
- Harris, W.V. (1967) Termites of the genus *Anacanthotermes* in North Africa and the near east (Isoptera: Hodotermitidae). *Proceedings of the Royal Entomological Society of London: Series B*, **36**, 79–86.
- Hausberger, B., Kimpel, D., van Neer, A. & Korb, J. (2011) Uncovering cryptic species diversity of a termite community in a West Africa savanna. *Molecular Phylogenetics and Evolution*, **61**, 964–969.
- Hebert, P.D.N., Cywinski, A., Ball, S.L. & DeWaard, J.R. (2003) Biological identifications through DNA barcodes. *Proceedings of the Royal Society B: Biological Sciences*, **270**, 313–321.
- Hellemans, S., Sobotnik, J., Lepoint, G., Mihaljevic, M., Roisin, Y. & Bourguignon, T. (2022) Termite dispersal is influenced by their diet. *Proceedings of the Royal Society B*, **289**, 20220246.
- Henschel, J.R. (1994) Diet and foraging behaviour of huntsman spiders in the Namib dunes (Araneae: Heteropodidae). *Journal of Zoology*, **234**, 239–251.
- Henschel, J.R. (1997) Psammophily in Namib desert spiders. *Journal of Arid Environments*, **37**, 695–707.
- Henschel, J.R. & Lubin, Y.D. (1992) Environmental factors affecting the web and activity of a psammophilous spider in the Namib Desert. *Journal of Arid Environments*, **22**, 173–189.
- Herzig-Straschil, B. & Herzig, A. (1989) Biology of *Xerus princeps* (Rodentia, Sciuridae). *Madoqua*, **16**, 41–46.
- Hibbitts, T.J., Whiting, M.J. & Stuart-Fox, D.M. (2007) Shouting the odds: vocalization signals status in a lizard. *Behavioral Ecology and Sociobiology*, **61**, 1169–1176.
- Holt, J.A. & Lepage, M. (2000) Termites and soil properties. *Termites: evolution, sociality, symbioses, ecology* (ed. by T. Abe, D.E. Bignell and M. Higashi), pp. 389–407. Springer Science & Business Media, Dordrecht, The Netherlands.
- Inward, D.J.G., Vogler, A.P. & Eggleton, P. (2007) A comprehensive phylogenetic analysis of termites (Isoptera) illuminates key aspects of their evolutionary biology. *Molecular Phylogenetics and Evolution*, **44**, 953–967.
- IUSS Working Group WRB (2015) World Reference Base for Soil Resources 2014, update 2015. International soil classification system for naming soils and creating legends for soil maps. *World Soil Resources Reports No. 106*. FAO, Rome.
- Izenman, A.J. (2008) *Modern multivariate statistical techniques. Regression, classification and manifold learning*. Springer, New York.
- Jacobson, K.M. (1996) Macrofungal ecology in the Namib Desert. A fruitful or futile study? *McIlvainea* 21–32.
- Jacobsen, N.H.G. & Moss, H. (1987) A contribution to the flora of the northern Namib. *Dinteria*, **19**, 27–68.
- Jacobson, K., van Diepeningen, A., Evans, S., Fritts, R., Gemmel, P., Marsho, C., Seely, M., Wenndt, A., Yang, X. & Jacobson, P. (2015) Non-rainfall moisture activates fungal decomposition of surface litter in the Namib Sand Sea. *PLoS One*, **10**, e0126977.
- Jankowitz, W.J., Van Rooyen, M.W., Shaw, D., Kaumba, J.S. & Van Rooyen, N. (2008) Mysterious circles in the Namib Desert. *South African Journal of Botany*, **74**, 332–334.
- Jones, C.G., Lawton, J.H. & Shachak, M. (1994) Organisms as ecosystem engineers. *Oikos*, **69**, 373–386.
- Jones, J.A. (1990) Termites, soil fertility and carbon cycling in dry tropical Africa: a hypothesis. *Journal of Tropical Ecology*, **6**, 291–305.
- Jouquet, P., Bottinelli, N., Shanbhag, R. R., Bourguignon, T., Traoré, S. & Abbasi, S. A. (2016) Termites: the neglected soil engineers of tropical soils. *Soil Science*, **181**, 157–165.
- Jouquet, P., Traoré, S., Choosai, C., Hartmann, C. & Bignell, D. (2011) Influence of termites on ecosystem functioning. Ecosystem services provided by termites. *European Journal of Soil Biology*, **47**, 215–222. <https://doi.org/10.1016/j.ejsobi.2011.05.005>
- Jouquet, P., Traoré, S., Harit, A., Choosai, C., Cheik, S. & Bottinelli, N. (2020) Moving beyond the distinction between the bright and dark sides of termites to achieve sustainable development goals. *Current Opinion in Insect Science*, **40**, 71–76. <https://doi.org/10.1016/j.cois.2020.05.010>
- Jürgens, N. (1986) Untersuchungen zur Ökologie sukkulenter Pflanzen des südlichen Afrika. *Mitteilungen aus dem Institut für Allgemeine Botanik Hamburg*, **21**, 139–365.
- Jürgens, N. (1991) A new approach to the Namib Region. *Vegetatio*, **97**, 21–38.
- Jürgens, N. (1996) Psammophorous plants and other adaptations to desert ecosystems with high incidence of sandstorms. *Feddes Repertorium*, **107**, 345–359.
- Jürgens, N. (1997) Floristic biodiversity and history of African arid regions. *Biodiversity & Conservation*, **6**, 495–514.
- Jürgens, N. (2004) A first classification of the vegetation of the Richtersveld (RSA) and directly adjacent regions in in Namibia and South Africa. *Biodiversity & Ecology*, **2**, 149–180.
- Jürgens, N. (2013) The biological underpinnings of Namib Desert fairy circles. *Science*, **339**, 1618–1621.
- Jürgens, N. (2015) Exploring common ground for different hypotheses on Namib fairy circles. *Ecography*, **38**, 12–14.
- Jürgens, N., Desmet, P.G., Rutherford, M.C., Mucina, L. & Ward, R.A. (2005) Desert Biome. *The Vegetation of South Africa, Lesotho and Swaziland* (ed. by L. Mucina and M.C. Rutherford), pp. 300–323. South African National Biodiversity Institute, Pretoria.
- Jürgens, N., Gunter, F., Oldeland, J., Gröngröft, A., Henschel, J.R., Oncken, I. & Picker, M.D. (2021b) Largest on earth: Discovery of a new type of fairy circle in Angola supports a termite origin. *Ecological Entomology*, **46**, 777–789.
- Jürgens, N., Haarmeyer, D.H., Luther-Mosebach, J., Dengler, J., Finckh, M. & Schmiedel, U. (2010) *Patterns at local scale: The BIOTA Observatories*. Klaus Hess Publishers, Göttingen & Windhoek.
- Jürgens, N., Oldeland, J., Hachfeld, B., Erb, E. & Schultz, C. (2013) Ecology and spatial patterns of large-scale vegetation units within the central Namib Desert. *Journal of Arid Environments*, **93**, 59–79.

- Jürgens, N., Oncken, I., Oldeland, J., Gunter, F. & Rudolph, B. (2021a) Welwitschia: Phylogeography of a living fossil, diversified within a desert refuge. *Scientific reports*, **11**, 1–14.
- Jürgens, N., Strohbach, B., Lages, F., Schmiedel, U., Finckh, M., Sichone, P., Hahn, L.M. & Zigelksi, P. (2018) Biodiversity observation—an overview of the current state and first results of biodiversity monitoring studies. *Biodiversity & Ecology*, **6**, 382–396.
- Jürgens, N., Vlieghe, K.E.P., Bohn, C., Erni, B., Gunter, F., Oldeland, J., Rudolph, B. & Picker, M.D. (2015) Weaknesses in the plant competition hypothesis for fairy circle formation and evidence supporting the sand termite hypothesis. *Ecological Entomology*, **40**, 661–668.
- Kaffarnik, S. (2009) GC/MS-Analytik von Boden- und Termitennestmaterial aus Feenkreisen, Doctoral dissertation, University of Hamburg.
- Kappel, C., Illing, N., Huu, C.N., Barger, N.N., Cramer, M.D., Lenhard, M. & Midgley, J.J. (2020) Fairy circles in Namibia are assembled from genetically distinct grasses. *Communications Biology*, **3**, 1–8.
- Kassambara, A. & Mundt, F. (2017) Factoextra: extract and visualize the results of multivariate data analyses. *R package version 1.0.5*. (accessed 8th November 2018).
- Klak, C., Hanacek, P. & Bruyns, P. (2017) Out of southern Africa: Origin, biogeography and age of the Aizooidae (Aizoaceae). *Molecular Phylogenetics and Evolution*, **109**, 203–216.
- Klak, C., Reeves, G. & Hedderson, T. (2004) Unmatched tempo of evolution in Southern African semi-desert ice plants. *Nature*, **427**, 63–65.
- Klingenberg, C.P. (2011) MorphoJ: an integrated software package for geometric morphometrics. *Molecular Ecology Resources*, **11**, 353–357.
- Kocarek, M. & Kodesova, R. (2012) Influence of temperature on soil water content measured by ECH2O-TE sensors. *International Agrophysics*, **26**, 259–269.
- Koch, C. (1962) The Tenebrionidae of Southern Africa, XXXI. Comprehensive notes on the tenebrionid fauna of the Namib Desert. *Annals of the Transvaal Museum*, **24**, 61–106.
- Kohfahl, C., Molano-Leno, L., Martínez, G., Vanderlinde, K., Guardiola-Albert, C. & Moreno, L. (2019) Determining groundwater recharge and vapor flow in dune sediments using a weighable precision meteo lysimeter. *Science of the Total Environment*, **656**, 550–557.
- Korb, J. (2003) Thermoregulation and ventilation of termite mounds. *Naturwissenschaften*, **90**, 212–219.
- Krasulová, J., Hanus, R., Kutalová, K., Šobotník, J., Sillam-Dussès, D., Tichý, M. & Valterová, I. (2012) Chemistry and Anatomy of the Frontal Gland in Soldiers of the Sand Termite *Psammotermes hybostoma*. *Journal of Chemical Ecology*, **38**, 557–565.
- Kuhn M. (2018) Caret: classification and regression training. *Astrophysics Source Code Library*, ascl-1505, R package version 6.0-80 (accessed 8th November 2018).
- Kuhn, M. & Johnson, K. (2013) *Applied predictive modeling*, 26th edn. Springer, New York.
- Kutner, M.H., Nachtsheim, C.J., Neter, J. & Li, W. (2005) *Applied Linear Statistical Models*. McGraw Hill Irwin, New York.
- Lancaster, N. (2014) Dune systems of the Namib Desert—a spatial and temporal perspective. *Transactions of the Royal Society of South Africa*, **69**, 133–137.
- Lasser, J., Nield, J.M., Ernst, M., Karius, V., Wiggs, G.F.S. & Goehring, L. (2019) Salt Polygons are caused by convection. *arXiv preprint arXiv*, **1902.03600**.
- Lê, S., Josse, J. & Husson, F. (2008) FactoMineR: An R package for multivariate analysis. *Journal of Statistical Software*, **25**, 1–18.
- Lefever, R. & Lejeune, O. (1997) On the origin of tiger bush. *Bulletin of Mathematical Biology*, **59**, 263–294.
- Legendre, P. & Gallagher, E.D. (2001) Ecologically meaningful transformations for ordination of species data. *Oecologia*, **129**, 271–280.
- Legendre, P. & Legendre, L. (2012) *Numerical ecology*, 24th edn. Developments in Environmental Modelling. Elsevier, Amsterdam.
- Lemke, K. (2018) A general study about the ant species *Carebara kunensis* of the Namib Desert with special regard to their foraging behaviour and activity. Bachelor thesis, Institute of Plant Science and Microbiology, University of Hamburg.
- Lewis, R.J. (2000) An introduction to classification and regression tree (CART) analysis. *Annual meeting of the society for academic emergency medicine in San Francisco, California*, Vol. 14, Citeseer, 1–14.
- Lewis, E.R., Narins, P.M., Jarvis, J.U., Bronner, G., & Mason, M.J. (2006) Preliminary evidence for the use of microseismic cues for navigation by the Namib golden mole. *The Journal of the Acoustical Society of America*, **119**, 1260–1268.
- Li, Y., Wang, Y. & Yao, Q. (2020) Nebkhas origination in arid and semi-arid regions: An overview. *Acta Ecologica Sinica*, **40**, 500–505.
- Lo, N., Kitade, O., Miura, T., Constantino, R. & Matsumoto, T. (2004). Molecular phylogeny of the Rhinotermitidae. *Insectes Sociaux*, **51**, 365–371.
- Loh, W.-Y. (2011) Classification and regression trees. *Wiley Interdisciplinary Reviews: Computational Statistics*, **1**, 14–23.
- Lorenz, C., Almeida, F., Almeida-Lopes, F., Louise, C., Pereira, S.N., Petersen, V., Vidal, P.O., Virdinio, F. & Suesdek, L. (2017) Geometric morphometrics in mosquitoes: What has been measured? *Infection, Genetics and Evolution*, **54**, 205–215.
- Louw, G.N. & Seely M.K. (1982) Ecology of desert organisms, 1st edn. Longman Group Limited, London and New York.
- Lovegrove, B.G. & Siegfried, W.R. (1986) Distribution and formation of Mima-like earth mounds in the western Cape Province of South Africa. *South African Journal of Science*, **82**, 432–436.
- Lubin, Y.D. & Henschel, J.R. (1990) Foraging at the thermal limit: burrowing spiders (*Seothyra*, Eresidae) in the Namib desert dunes. *Oecologia*, **84**, 461–467.
- Lüscher, M. (1961) Air-conditioned termite nests. *Scientific American*, **205**, 138–145.
- Macfadyen W.A. (1950) Vegetation patterns in the semi desert plains of the British Somaliland. *Geographic Journal*, **116**, 199–210.
- Maestre, F.T., Benito, B.M., Berdugo, M., Concostrina-Zubiri, L., Delgado-Baquerizo, M., Eldridge, D.J., Guirado, E., Gross, N., Kéfi, S., Le Bagousse-Pinguet, Y., Ochoa-Hueso, R. & Soliveres, S. (2021) Biogeography of global drylands. *New Phytologist*, **231**, 540–558.
- Maher, B.A. (1998) Magnetite biominerilization in termites. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, **265**, 733–737.
- Marsh, A. C. (1987) Thermal responses and temperature tolerance of a desert ant-lion larva. *Journal of Thermal Biology*, **12**, 295–300.
- Martin, M.M. & Martin J.S. (1978.) Cellulose digestion in the midgut of the fungus-growing termite *Macrotermes natalensis*: the role of acquired digestive enzymes. *Science*, **199**, 1453–1455.
- Matimati, I., Musil, C.F., Raitt, L. & February, E. (2013) Non rainfall moisture interception by dwarf succulents and their relative abundance in an inland arid South African ecosystem. *Ecohydrology*, **6**, 818–825.
- McAuliffe, J.R., Hoffman, M.T., McFadden, L.D., Bell, W., Jack, S., King, M.P. & Nixon, V. (2019a) Landscape patterning created by the southern harvester termite, *Microhodotermes viator*: Spatial dispersion of colonies and alteration of soils. *Journal of Arid Environments*, **162**, 26–34.
- McAuliffe, J.R., Hoffman, M.T., McFadden, L.D., Jack, S., Bell, W. & King, M.P. (2019b) Whether or not heuweltjies: context-dependent ecosystem engineering by the southern harvester termite, *Microhodotermes viator*. *Journal of Arid Environments*, **163**, 26–33.
- McAuliffe, J.R., Hoffman, T.M., McFadden, L.D. & King, M.P. (2014) Role of aeolian sediment accretion in the formation of heuweltjie earth mounds, western South Africa. *Earth Surface Processes and Landforms*, **39**, 1900–1912.
- Mendelsohn J. & Mendelsohn, S. (2018) Sudoeste de Angola: um Retrato da Terra e da Vida. South West Angola: a portrait of land and life. Arte e Ciência.

- Meron, E. (2012) Pattern-formation approach to modelling spatially extended ecosystems. *Eco-logical Modelling*, **234**, 70–82.
- Meron, E. (2015) *Nonlinear Physics of Ecosystems*. CRC Press Boca Raton. 329 pp.
- Meron, E. (2019) Vegetation pattern formation: The mechanisms behind the forms. *Physics Today*, **72**, 30–36. <https://doi.org/10.1063/PT.3.4340>
- Meyer, J.J.M., Heyman, H.M., Senejoux, F., Degashu, P., Galt, N., Nxumalo, D. & Labuschagne, N. (2017) The chemical ‘footprints’ of *Euphorbia* species found in the mysterious fairy circles of Namibia. *South African Journal of Botany*, **100**, 351.
- Meyer, J.M., Schutte, C.S., Galt, N., Hurter, J.W. & Meyer, N.L. (2021) The fairy circles (circular barren patches) of the Namib Desert—What do we know about their cause 50 years after their first description? *South African Journal of Botany*, **140**, 226–239.
- Meyer, J.J.M., Schutte, C.E., Hurter, J.W., Galt, N.S., Degashu, P., Breetzke, G., Baranenko, D. & Meyer, N. L. (2020) The allelopathic, adhesive, hydrophobic and toxic latex of *Euphorbia* species is the cause of fairy circles investigated at several locations in Namibia. *BMC ecology*, **20**, 1–23.
- Meyer, J.J.M., Senejoux, F., Heyman, H.M., Meyer, N.L. & Meyer, M.A. (2015) The occurrence of triterpenoids from *Euphorbia gummifera* inside the fairy circles of Garub in the southern Namibian pro-desert. *South African Journal of Botany*, **98**, 10–15.
- Meyer, V.W., Braack, L.E.O., Biggs, H.C. & Ebersohn, C. (1999) Distribution and density of termite mounds in the northern Kruger National Park, with specific reference to those constructed by *Macrotermes Holmgren* (Isoptera: Termitidae). *African Entomology*, **7**, 123–130.
- Mills, A.J., Milewski, A., Fey, M.V., Groengroef, A. & Petersen, A. (2009) Fungus culturing, nutrient mining and geophagy: a geochemical investigation of *Macrotermes* and *Trinervitermes* mounds in southern Africa. *Journal of Zoology*, **278**, 24–35.
- Möhlig, W. (2002) The language history of Herero as a source of ethnohistorical interpretations. *People, Cattle and Land: Transformations of a pastoral society in southwestern Africa* (ed. by M. Bollig and J.-B. Gewald), pp. 119–146. Rüdiger Köppe, Köln.
- Moll, E.J. (1994) The origin and distribution of fairy rings in Namibia. *Proceedings of the 13th Plenary Meeting AETFAT*, Malawi, **2**, 1203–1209.
- Moore, J.M. & Picker, M.D. (1991) Heuweltjies (earth mounds) in the Clanwilliam District, Cape Province, South Africa: 4000-year-old termite nests. *Oecologia*, **86**, 424–432.
- Muche, G., Schmiedel, U. & Jürgens, N. (2012) BIOTA Southern Africa biodiversity observatories vegetation database. *Environmental Monitoring and Assessment*, **184**, 655–678.
- Muche, G., Schmiedel, U., Finckh, M. & Jürgens, N. (2018) BIOTABase – a unique software to handle complex biodiversity observation data and environmental data. *Climate change and adaptive land management in southern Africa – Assessments, Changes, Challenges and Solutions* (ed. by R. Revermann, K. Krewnka, U. Schmiedel, J.M. Olwoch, J. Helmschrot, and N. Jürgens), pp. 458–459. *Biodiversity & Ecology*, 6, Klaus Hess Publishers, Göttingen & Windhoek.
- Mucina, L., Jürgens, N., Le Roux, A., Rutherford, M.C., Schmiedel, U., Esler, K.J., Powrie, L.W., Desmet, P.G. & Milton, S.J. (2006) Succulent Karoo Biome. *Strelitzia*, **19**, 220–299.
- Mühlenpfordt, M. (2013) Activity patterns and foraging behavior of *Messor denticornis* ants in fairy circle ecosystems of the Namib Desert. Bachelor thesis, Institute of Plant Science and Microbiology, University of Hamburg.
- Müller, B., Linstädter, A., Frank, K., Bollig, M. & Wissel, C. (2007) Learning from local knowledge: modeling the pastoral-nomadic range management of the Himba, Namibia. *Ecological applications*, **17**, 1857–1875.
- Myagkiy, A., Moretti, I. & Brunet, F. (2020) Space and time distribution of subsurface H₂ concentration in so-called “fairy circles”: Insight from a conceptual 2-D transport. *Bulletin de la Société Géologique de France-Earth Sciences Bulletin*, **191**, 13.
- Naiman, R.J., Melillo, J.M. & Hobbie, J.E. (1986) Ecosystem alteration of boreal forest streams by beaver (*Castor canadensis*). *Ecology*, **67**, 1254–1269.
- Naude, Y., Van Rooyen, M.W. & Rohwer, E.R. (2011) Evidence for a geochemical origin of the mysterious circles in the Pro-Namib desert. *Journal of Arid Environments*, **75**, 446–456.
- Noirot, C. & Darlington, J.P.E.C. (2000) Termite nests: Architecture, regulation and defence. *Termites: evolution, sociality, symbioses, ecology* (ed. by T. Abe, D.E. Bignell and M. Higashi), pp. 121–139. Springer Netherlands, Dordrecht.
- Oguz, I., Stocker, B. & Jürgens, N. (2004) Succulent plant communities and edaphical factors along a coast-inland-transect in the Sandveld of the Northern Richtersveld. *Biodiversity & Ecology*, **2**, 133–149.
- Ollier, C.D. & Seely, M. (1977) Patterned ground near Gobabeb, central Namib desert. *Journal of Arid Environments*, **10**, 213–214.
- Or, D. & Wraith, J.M. (1999) Temperature effects on soil bulk dielectric permittivity measured by time domain reflectometry: A physical model. *Water Resources Research*, **35**, 371–383.
- Orlóci, L. (1967) An agglomerative method for classification of plant communities. *The Journal of Ecology*, **55**, 193–206.
- Patton, P.C., Pickup, G. & Price, D.M. (1993) Holocene palaeofloods of the Ross River, central Australia. *Quaternary Research*, **40**, 201–212.
- Phan, T.H., Van Gorp, H., Li, Z., Huynh, T.M.T., Fujita, Y., Verstraete, L., Eyley, S., Thielemans, W., Uji-I, H., Hirsch, E.B., Mertens, S.F.L., Greenwood, J., Ivasenko, O. & De Feyter, S. (2019) Graphite and graphene fairy circles: A bottom-up approach for the formation of nanocorals. *ACS nano*, **13**, 5559–5571.
- Picker, M.D., Hoffman, M.T. & Leverton, B. (2007) Density of *Microhodotermes viator* (Hodotermitidae) mounds in southern Africa in relation to rainfall and vegetative productivity gradients. *Journal of Zoology*, **271**, 37–44.
- Picker, M.D., Ross-Gillespie, V., Vlieghe, K. & Moll, E. (2012) Ants and the enigmatic Namibian fairy circles—cause and effect? *Ecological Entomology*, **37**, 33–42.
- Pickford, M., Senut, B. & Dauphin, Y. (1995) Biostratigraphy of the Tsondab sandstone (Namibia) based on gigantic avian eggshells. *Geobios*, **28**, 85–98.
- Platnick, N.I. & Griffin, E. (1990) On *Rastellus*, a new genus of the spider family Ammoxenidae (Araneae, Gnaphosoidea). *American Museum Novitates*, **2995**.
- Potts, A.J., Midgley, J.J. & Harris, C. (2009) Stable isotope and 14C study of biogenic calcrete in a termite mound, Western Cape, South Africa, and its palaeoenvironmental significance. *Quaternary Research*, **72**, 258–264.
- Pringle, R.M. & Tarnita, C.E. (2017) Spatial self-organization of ecosystems: integrating multiple mechanisms of regular-pattern formation. *Annual Review of Entomology*, **62**, 359–377.
- Prötz, D., Heß, M., Schwager, M., Glaw, F. & Scherz, M.D. (2021) Neon-green fluorescence in the desert gecko *Pachydactylus rangei* caused by iridophores. *Scientific Reports*, **11**, 1–10.
- Ramond, J., Pienaar, A., Armstrong, A., Seely, M. & Cowan, D.A. (2014) Niche-Partitioning of Edaphic Microbial Communities in the Namib Desert Gravel Plain Fairy Circles. *PLoS ONE*, **9**, e109539. doi:10.1371/journal.pone.0109539
- Ravi, S., D’Odorico, P., Wang, L. & Collins, S. (2008) Form and function of grass ring patterns in arid grasslands: the role of abiotic controls. *Oecologia*, **158**, 545–555.
- Ravi, S., Wang, L., Kaseke, K.F., Buynevich, I.V. & Marais, E. (2017) Ecohydrological interactions within “fairy circles” in the Namib Desert: Revisiting the self-organization hypothesis. *Journal of Geophysical Research: Biogeosciences*, **122**, 405–414.
- Renger, M., Bohne, K., Facklam, M., Harrach, T., Rick, W., Schäfer, W., Wessolek, G. & Zacharias, S. (2009) Ergebnisse und Vorschläge der DBG-Arbeitsgruppe „Kennwerte des Bodengefüges“ zur Schätzung bodenphysikalischer Kennwerte. *Wilhadi*, **5**, 21682.

- Revermann, R., Krewenka, K., Schmiedel, U., Olwoch, J.M., Helmschrot, J. & Jürgens, N. (eds.) (2018) Climate change and adaptive land management in southern Africa – Assessments, Changes, Challenges and Solutions. Klaus Hess Publishers, Göttingen & Windhoek.
- Rietkerk, M. & van de Koppel, J. (2008) Regular pattern formation in real ecosystems. *Trends in Ecology and Evolution*, **23**, 169–175.
- Robinson, E.J. (2014) Polydomy: the organisation and adaptive function of complex nest systems in ants. *Current Opinion in Insect Science*, **5**, 37–43.
- Rodell, M., Houser, P.R., Jambor, U.E.A., Gottschalck, J., Mitchell, K., Meng, C. J., Arseneault, K., Cosgrove, B., Radakovich, J., Bosilovich, M., Entin, J. K., Walker, J. P., Lohmann, D. & Toll, D. (2004) The global land data assimilation system. *Bulletin of the American Meteorological Society*, **85**, 381–394.
- Rohlf, F.J. (2017) *tpsDig version 2.31*. New York: Department of Ecology and Evolution and Anthropology, State University of New York at Stony Brook.
- Rohlf, F.J. (2018) *tpsUtil version 1.75*. New York: Department of Ecology and Evolution and Anthropology, State University of New York at Stony Brook.
- Roisin, Y. & Lenz, M. (1999) Caste developmental pathways in colonies of *Coptotermes lacteus* (Froggatt) headed by primary reproductives (Isoptera, Rhinotermitidae). *Insectes Sociaux*, **46**, 273–280.
- Roonwal, M.L. & Bose, G. (1962) An African genus *Psammotermes* in Indian termite fauna, with fuller description of *P. rajasthanicus* from Rajasthan, India. *Records Indian Museum*, **58**, 151–158.
- Roonwal, M.L. & Chhotani, O.B. (1989) Fauna of India, Isoptera (Termites). *Zoological Survey of India*.
- Roy, V., Demanche, C., Livet, A. & Harry, M. (2006) Genetic differentiation in the soil-feeding termite *Cubitermes sp. affinis subarquatus*: occurrence of cryptic species revealed by nuclear and mitochondrial markers. *BMC Evolutionary Biology*, **6**, 102. <https://doi.org/10.1186/1471-2148-6-102>.
- Ruiz-Reynés, D., Gomila, D., Sintes, T., Hernández-García, E., Marbà, N. & Duarte, C. M. (2017) Fairy circle landscapes under the sea. *Science Advances*, **3**, e1603262.
- Rutherford, M.C. & Westfall, R.H. (1994) *Biomes of southern Africa: an objective categorization*. National Botanical Institute, Pretoria.
- Sanabria, C., Dubs, F., Lavelle, P., Fonte, S.J. & Barot, S. (2016) Influence of regions, land uses and soil properties on termite and ant communities in agricultural landscapes of the Colombian Llanos. *European Journal of Soil Biology*, **74**, 81–92.
- Sasakiwa, K. (2016) Utility of geometric morphometrics for inferring feeding habit from mouthpart morphology in insects: Tests with larval Carabidae (Insecta: Coleoptera). *Biological Journal of the Linnean Society*, **118**, 394–409.
- Schmiedel, D.A., Benítez, H.A., Borissov, I.M., & Fruciano, C. (2015) Bat species comparisons based on external morphology: a test of traditional versus geometric morphometric approaches. *PloS one*, **10**, e0127043.
- Schutte, C.E. (2019) Predictive modelling of Namibian fairy circles and their co-occurrence with *Euphorbia* species using Geographical Information Systems (GIS). Doctoral Dissertation, University of Pretoria.
- Schutte, C.E., Meyer, J.J.M. & Breetzke, G. (2018) Are the fairy circles of Namibia caused by *Euphorbia* species? *South African Journal of Botany*, **115**, 308.
- Shaanika, H.N. (2020) Assessing the likelihood that burrowing gerbils in the central Namib are ecological engineers. Namibia University of Science and Technology, Windhoek.
- Shaanika, H., Wassenaar, T. & Eiseb, S. (2019) *Gerbils of the Namib Desert—are they ecological engineers?* GOBABEB Namib Research Institute, Gobabeb.
- Sheffer, E., Yizhaq, H., Gilad, E., Shachak, M. & Meron, E. (2007) Why do plants in resource-deprived environments form rings? *Ecological Complexity*, **4**, 192–200.
- Sieben, A.J., Barger, N.N., Cramer, M.D., Tschinkel, W.R., Wiegand, T. & Shoemaker, L.G. (2019) Herbivory and soil resources impact the maintenance of landscape gap patterning and the spatial arrangement of grasses during pattern collapse. *ESA Annual Meeting (August 11–16)*. ESA.
- Sileshi, G.W., Arshad, M.A., Konaté, S. & Nkunku, P.O. (2010) Termite-induced heterogeneity in African savanna vegetation: mechanisms and patterns. *Journal of Vegetation Science*, **21**, 923–937.
- Sillam-Dussès, D., Hanus, R., Abd El-Latif, A. O., Jiroš, P., Krasulová, J., Kalinová, B., Valterová, I. & Šobotník, J. (2011) Sex Pheromone and Trail Pheromone of the Sand Termite *Psammotermes hybostoma*. *Journal of Chemical Ecology*, **37**, 179–188.
- Silvestri, F. (1908) B. Archiptera (I.). Termitidae. *Denkschriften der Medicinisch-Naturwissenschaftlichen Gesellschaft zu Jena*, **13**, 69–82.
- Sjöstedt, Y. (1900) Monographie der Termiten Afrikas. *Kungliga Svenska Vetenskapsakademiens Handlingar*, **34**, 1–236.
- Sjöstedt, Y. (1912) Neue Termiten aus Tripolis, Ober-Ägypten, Abessinien, Eritrea, dem Gallaland und Somaliland. *Wiksell's Boktryckeri*, Uppsala.
- Sjöstedt, Y. (1925) Revision der Termiten Afrikas 3. Monographie. *Kungliga Svenska Vetenskapsakademiens Handlingar*, **3**, 1–419.
- Skiljan, I. (2012). „IrfanView.“ URL <http://irfanview.tuwien.ac.at/>.
- Skinner, J.D. & Chimimba, C.T. (2013) *Mammals of the southern African subregion*. Cambridge University Press, Cambridge.
- Slavich, P.G. & Petterson, G.H. (1993) Estimating the electrical conductivity of saturated paste extracts from 1: 5 soil, water suspensions and texture. *Soil Research*, **31**, 73–81.
- Sokol, J. (2018) Visiting the Mysterious Fairy Circles of the Namib Desert. *The Atlantic* <https://www.theatlantic.com/science/archive/2018/05/visiting-the-mysterious-fairy-circles-of-the-namib-desert/560639/>
- Spaargaren, O.C. & Deckers, J. (1998) The world reference base for soil resources. *Soils of Tropical Forest Ecosystems*. Springer.
- Stander, P.E. (2019) Lions (*Panthera leo*) specialising on a marine diet in the Skeleton Coast National Park, Namibia. *Namibian Journal of Environment*, **3**, 1–10.
- Stengel, I. (2000) Brousse tigré – Muster und Materialverspülung und Pflanzenwuchs in Südnamibia. *Petermanns Geographische Mitteilungen*, **144**(2), 4–5.
- Stronen, A.V., Navid, E.L., Quinn, M.S., Paquet, P.C., Bryan, H.M. & Darimont, C.T. (2014) Population genetic structure of gray wolves (*Canis lupus*) in a marine archipelago suggests island-mainland differentiation consistent with dietary niche. *BMC Ecology*, **14**, 1–9.
- Swart, R. (2008) How old is the Namib really – A geological perspective. *Namib: Secrets of a desert uncovered* (ed. by M.K. Seely and J. Pallett), pp. 1–201. Venture Publications, Windhoek.
- Tarnita, C.E., Bonachela, J.A., Sheffer, E., Guyton, J.A., Coverdale, T.C., Long, R.A. & Pringle, R.M. (2017) A theoretical foundation for multi-scale regular vegetation patterns. *Nature*, **541**, 398–401.
- Taylor, W.A., Lindsey, P.A. & Skinner, J.D. (2002) The feeding ecology of the aardvark *Orycteropus afer*. *Journal of Arid Environments*, **50**, 135–152.
- Tevis, L.J. & Newell, I.M. (1962) Studies on the biology and seasonal cycle of the giant red velvet mite, *Dinothrombium pandorae* (Acar, Trombidiidae). *Ecology*, **43**, 497–505.
- Therneau, T., Atkinson, B. & Ripley, B. (2018) rpart: Recursive Partitioning and Regression Trees. R package version 4.1–13.
- Theron, G.K. (1979) Die verskynsel van kaal kolle in Kaokoland, Suidwes-Afrika. *Journal of the South African Biological Society*, **20**, 43–53.
- Tinley, K.L. (1971) Etosha and the Kaokoveld. *African Wild Life Protection Society*, **25**, 1–16.
- Thidi, M., Lefever, R. & Vladimirov, A. (2008) On Vegetation Clustering, Localized Bare Soil Spots and Fairy Circles. *Lecture Notes in Physics*, **751**, 381–402. https://doi.org/10.1007/978-3-540-78217-9_15

- Tschinkel, W.R. (2010) The foraging tunnel system of the Namibian Desert termite, *Baicaliotermes hainesi*. *Journal of Insect Science*, **10**, 65, 1–17.
- Tschinkel, W.R. (2012) The life cycle and life span of Namibian fairy circles. *PLoS one*, **7**, e38056.
- Tschinkel, W.R. (2015) Blog to Ecography. Comments by WR Tschinkel in response to Ecography article by Juergens, N. <https://www.ecography.org/blog/new-studies-fairy-circles-need-account-new-observations-their-spatial-patterns> (accessed 14th August 2021).
- Turner, J.S. (2001) On the mound of *Macrotermes michaelseni* as an organ of respiratory gas exchange. *Physiological and Biochemical Zoology*, **74**, 798–822.
- Uys, V.M. (2002) *A guide to the termite genera of southern Africa* (No. 15). Plant Protection Research Institute, Agricultural Research Council, Pretoria.
- Van der Walt, A.J., Johnson, R.M., Cowan, D.A., Seely, M., Ramond, J. (2016) Unique microbial phylotypes in Namib Desert dune and gravel plain Fairy Circle soils. *Applied and Environmental Microbiology*, **82**, 4592–4601. <https://journals.asm.org/doi/10.1128/AEM.00844-16>
- Van Rooyen, M.W., Theron, G.K., Van Rooyen, N., Jankowitz, W.J. & Matthews, W.S. (2004) Mysterious circles in the Namib Desert: review of hypotheses on their origin. *Journal of Arid Environments*, **57**, 467–485.
- Viljoen, P.J. (1980) Veldtipies, verspreiding van die groter soogdiere en enkele aspekte van die ekologie van Kaokoland. Doctoral dissertation, University of Pretoria.
- Vlieghe, K. & Picker, M. (2019) Do high soil temperatures on Namibian fairy circle discs explain the absence of vegetation? *Plos one*, **14**, e0217153.
- Vlieghe, K., Picker, M.D., Ross-Gillespie, V. & Ernie, B. (2014) Herbivory by subterranean termite colonies and the development of fairy circles in SW Namibia. *Ecological Entomology*, **40**, 42–49.
- Wagner, T.C., Richter, J., Joubert, D.F. & Fischer, C. (2018) A dominance shift in arid savanna: An herbaceous legume outcompetes local C₄ grasses. *Ecology and Evolution*, **8**, 6779–6787.
- Wagner, T.C., Uiseb, K. & Fischer, C. (2021) Rolling pits of Hartmann's mountain zebra (*Zebra equus hartmannae*) increase vegetation diversity and landscape heterogeneity in the Pre-Namib. *Ecology and Evolution*, **11**, 13036–13051.
- Walsh, F.J., Sparrow, A.D., Kendrick, P. & Schofield, J. (2016) Fairy circles or ghosts of termitaria? Pavement termites as alternative causes of circular patterns in vegetation of desert Australia. *Proceedings of the National Academy of Sciences*, **113**, E5365–E5367.
- Walsh, F., Kendrick, P., Evans, T., Moore, D., Nelson, M., Oldham, C., Schofield, J. & Sparrow, A. (2022, preprint) First Peoples and scientists concur that 'fairy circles' are termite linyji, Research Square.
- Wan, T., Liu, Z., Leitch, I.J. et al. (2021) The *Welwitschia* genome reveals a unique biology underpinning extreme longevity in deserts. *Nature Communications*, **12**, 1–15.
- Wang, X. (2015) Vapor flow resistance of dry soil layer to soil water evaporation in arid environment: An overview. *Water*, **7**, 4552–4574.
- Ward, J.D. (1988) Eolian, fluvial and pan (playa) facies of the Tertiary Tsodab Sandstone Formation in the central Namib Desert, Namibia. *Sedimentary Geology*, **55**, 143–162.
- Ward, J.D. & Corbett, I. (1990) Towards an age of the Namib. *Namib Ecology*, **25**, 17–26.
- Ward, J.D., Seely, M.K. & Lancaster, N. (1983) On the antiquity of the Namib. *South African Journal of Science*, **79**, 175–183.
- Watson, A. (1980) Vegetation polygons in the central Namib Desert near Gobabeb. *Journal of Arid Environments*, **11**, 315–325.
- Watson, J.A.L. & Gay, F.J. (1970) The role of grass-eating termites in the degradation of a mulga ecosystem. *Search*, **1**, 1–43.
- Watson, J.A.L., Lendon, C. & Low, B.S. (1973) Termites in Mulga Lands. *Tropical Grasslands*, **7**, 121–126.
- Webb, J.K., Shine, R., Branch, W.R. & Harlow, P.S. (2000) Life history strategies in basal snakes: reproduction and dietary habits of the African thread snake, *Leptotyphlops scutifrons* (Serpentes, Leptotyphlopidae). *Journal of Zoology*, **250**, 321–327.
- Weesner, F. (1970) External Anatomy. *Biology of termites*, Vol. I (ed. by K. Krishna & F.M. Weesner), pp. 1–23. Academic Press, New York.
- Wells, S.G. & Dohrenwend, J.C. (1985) Relict sheetflood bed forms on late Quaternary alluvial-fan surfaces in the southwestern United States. *Geology*, **13**, 512–516.
- Wharton, R.A. (1981) Namibian Solifugae (Arachnida). *Cimbebasia Memoir*, **5**, 1–87.
- Whiteford, W.G. & Eldridge, D.J. (2013) Effects of ants and termites on soil and geomorphological processes. *Treatise on Geomorphology*, **12**, 281–292.
- Whittington-Jones, G.M., Bernard, R.T.F. & Parker, D.M. (2011) Aardvark burrows: a potential resource for animals in arid and semi-arid environments. *African Zoology*, **46**, 362–370.
- Wildermuth, B. (2020) Spatial patterns, life history and biological interactions of termite colonies (*Macrotermes* sp.) in the thorn bush savanna of Central Namibia. Master thesis, Institute of Plant Science and Microbiology, University of Hamburg.
- Wildermuth, B., Oldeland, J. & Juergens, N. (2021) A beneficial relationship: associated trees facilitate termite colonies (*Macrotermes michaelseni*) in Namibia. *Ecosphere*, **12**, e03671.
- Will, K.W., Mishler, B.D. & Wheeler, Q.D. (2005) The perils of DNA barcoding and the need for integrative taxonomy. *Systematic Biology*, **54**, 844–851.
- Williams, L.A.D., Vasques, E., Reid, W., Porter, R. & Kraus, W. (2003) Biological activities of an extract from *Cleome viscosa* L. (Capparaceae). *Naturwissenschaften*, **90**, 468–472.
- Williams, R.M.C. (1966) The east African termites of the genus *Cubitermes* (Isoptera: Termitidae). *Transactions of the Royal Entomological Society of London*, **118**, 73–118.
- Yizhaq, H., Al-Tawaha, A.R.M.S. & Stavi, I. (2022) A first study of *Urginea maritima* rings: A case study from southern Jordan. *Land*, **11**, 285. <http://dx.doi.org/10.3390/land11020285>.
- Zanne, A.E., Flores-Moreno, H., Powell, J.R. et al. (2022) Termite sensitivity to temperature affects global wood decay rates. *Science*, **377**, 1440–1444. <https://doi.org/10.1126/science.abo3856>
- Zelditch, M.L., Swiderski, D.L., Sheets, H.D. & Fink, W.L. (2004) *Geometric morphometrics for biologists*. Elsevier, Amsterdam.
- Zeidler, J. (1997) *Distribution of termites (Isoptera) throughout Namibia-environmental connections*. Doctoral dissertation, University of Witwatersrand.
- Zelnik, Y.R., Meron, E. & Bel, G. (2015) Gradual regime shifts in fairy circles. *Proceedings of the National Academy of Sciences*, **112**, 12327–12331.